arXiv:hep-ph/0512140v1 12 Dec 2005

Transversity[1]

Mauro Anselmino*

*Dipartimento di Fisica Teorica, Università di Torino and
INFN, Sezione di Torino, Via P. Giuria 1, I-10125 Torino, Italy

Abstract

Some general remarks on parton transverse spin distributions and transverse motions are presented. The issue of accessing experimental information on the transversity distributions h_1^q is discussed. In particular direct information could be obtained from Drell-Yan processes with polarized protons and antiprotons (planned at GSI), while indirect information, coupling h_1^q to the Collins fragmentation function, is being gathered by HERMES and COMPASS collaborations. The related issue of transverse Single Spin Asymmetries (SSA) measured in several processes and the role of the Sivers distribution function is also discussed.

1. Towards transversity distributions: theory and experiments

The partonic structure of protons and neutrons is well known concerning the longitudinal degrees of freedom; these refer to the longitudinal momentum and spin carried by the partons inside unpolarized or longitudinally polarized fast moving nucleons. Instead, much less is known about the transverse – with respect to the direction of motion – degrees of freedom: the intrinsic motion of partons ($\boldsymbol{k}_\perp$) and the transverse spin distributions of quarks inside a transversely polarized nucleon (transversity). Their knowledge – including possible spin-$\boldsymbol{k}_\perp$ correlations – is crucial if we aim at having a full understanding of the nucleon structure in terms of spin and orbital motion of quarks and gluons.

Transversity is the last leading-twist missing information on the quark spin structure of the nucleon [1]; whereas the unpolarized quark distributions, $q(x, Q^2)$ or $f_{q/p}(x, Q^2)$, are very well known, and good information is now available on the quark helicity distributions $\Delta q(x, Q^2)$, nothing is experimentally known on the nucleon transversity distribution $h_1^q(x, Q^2)$ [also denoted by $\Delta_T q(x, Q^2)$ or $\delta q(x, Q^2)$]. The reason why h_1^q, despite its fundamental importance, has never been measured is that it is a chiral-odd function, and consequently it decouples from inclusive Deep

---

[1] Summary of talks delivered at the Workshop "Hadron Structure and Hadron Spectroscopy", Prague, August 1-3, 2005 and at the International Workshop on "Transverse Polarisation Phenomena in Hard Processes" (Transversity 2005), Como, September 7-10, 2005.

Inelastic Scattering (DIS), which is our usual main source of information on the nucleon partonic structure. Since electroweak and strong interactions conserve chirality, h_1^q cannot occur alone, but has to be coupled to a second chiral-odd quantity.

This is possible, for example, in polarized Drell-Yan processes [2], where one measures the product of two transversity distributions, and in semi-inclusive DIS, where one couples h_1^q to a new unknown, chiral-odd, fragmentation function, the so-called Collins function [3].

1.1 h_1^q in Drell-Yan processes

Measurement of transversity is planned at RHIC, in Drell-Yan processes with transversely polarized protons, $p^\uparrow p^\uparrow \to \ell^- \ell^+ X$, via the measurement of the double spin asymmetry:

$$A_{TT} \equiv \frac{d\sigma^{\uparrow\uparrow} - d\sigma^{\uparrow\downarrow}}{d\sigma^{\uparrow\uparrow} + d\sigma^{\uparrow\downarrow}}, \tag{1}$$

which reads at leading order in the parton model

$$A_{TT}^{pp} = \hat{a}_{_{TT}} \, \frac{\sum_q e_q^2 \left[h_1^q(x_1, M^2)\, h_1^{\bar{q}}(x_2, M^2) + h_1^{\bar{q}}(x_1, M^2)\, h_1^q(x_2, M^2) \right]}{\sum_q e_q^2 \left[q(x_1, M^2)\, \bar{q}(x_2, M^2) + \bar{q}(x_1, M^2)\, q(x_2, M^2) \right]} \,, \tag{2}$$

where $q = u, d, s$; M is the invariant mass of the lepton pair and $\hat{a}_{TT}$ is the double spin asymmetry of the QED elementary process, $q\,\bar{q} \to \ell^- \ell^+$ [see below, Eq. (6)]. In this case one measures the product of two transversity distributions, one for a quark and one for an anti-quark. The latter (in a proton) is expected to be small; moreover, the QCD evolution of transversity is such that, in the kinematical regions of RHIC data, $h_1^q(x, Q^2)$ is much smaller than the corresponding values of $\Delta q(x, Q^2)$ and $q(x, Q^2)$. All this makes the Drell-Yan double spin asymmetry A_{TT}^{pp} expected at RHIC very small, no more than a few percents [4, 5].

Definite and direct information on transversity could be best obtained by measuring the double transverse spin asymmetry A_{TT} in the collision of polarized protons and antiprotons. The kinematical regions should be such that $h_1^q(x, Q^2)$ is expected to be sizeable, and the cross section allows to reach a good statistics. Such an ideal situation can be obtained by having polarized anti-protons colliding on polarized protons in the High Energy Storage Ring at GSI, as proposed by the PAX collaboration [6]. The expected ranges of energy and M^2 are:

$$(30 \lesssim s \lesssim 210)\ \mathrm{GeV}^2; \qquad M \gtrsim 2\ \mathrm{GeV}/c^2; \qquad \tau = x_1 x_2 = \frac{M^2}{s} \gtrsim 0.02\,. \tag{3}$$

One has

$$\begin{aligned} A_{TT}^{p\bar{p}} &= \hat{a}_{_{TT}} \, \frac{\sum_q e_q^2 \left[h_1^q(x_1)\, h_1^q(x_2) + h_1^{\bar{q}}(x_1)\, h_1^{\bar{q}}(x_2) \right]}{\sum_q e_q^2 \left[q(x_1)\, q(x_2) + \bar{q}(x_1)\, \bar{q}(x_2) \right]} & (4) \\ &\simeq \hat{a}_{_{TT}} \, \frac{h_1^u(x_1)\, h_1^u(x_2)}{u(x_1)\, u(x_2)} \quad \text{(at large } x) \,, & (5) \end{aligned}$$

where all quark distributions refer to protons and $\hat{a}_{TT}$ is the elementary double spin asymmetry for the $q\,\bar{q} \to \ell^- \ell^+$ process (see, *e.g.*, Ref. [1] for the definition of the scattering angles):

$$\hat{a}_{TT}(\theta, \varphi) = \frac{\sin^2\theta}{1+\cos^2\theta}\,\cos(2\varphi)\,. \tag{6}$$

Predictions [7, 8], based on assuming at some initial scale $h_1^q = \Delta q$ or on a chiral quark-soliton model, indicate large expected values of $A_{TT}^{p\bar{p}}$. Also the QCD corrections have been computed [9]; although they are huge for the cross sections, they affect very little the asymmetry, as shown in Fig. 1.



Figure 1: Expected values of $A_{TT}^{p\bar{p}}$ for the proposed PAX-GSI experiments with polarized proton and antiprotons. The figure is taken from Ref. [9]: $s = 210$ (GeV)^2, the transversity distributions are given by saturation of the Soffer bound, $2|h_1^q(x)| = q(x) + \Delta q(x)$, and higher order perturbative QCD corrections have been computed, including all-order soft-gluon resummations

1.2 h_1 from SIDIS processes and extraction of the Collins function

The Collins mechanism describes the fragmentation function of a transversely polarized quark (with spin vector $\boldsymbol{S}_q$) into an unpolarized or spinless hadron h (typically a pion):

$$\hat{D}_{h/q^\uparrow}(z, \boldsymbol{p}_\perp) = \hat{D}_{h/q}(z, p_\perp) + \frac{1}{2}\,\Delta^N\hat{D}_{h/q^\uparrow}(z, p_\perp)\,\boldsymbol{S}_q \cdot (\hat{\boldsymbol{p}}_q \times \hat{\boldsymbol{p}}_\perp)\,, \tag{7}$$

where $\boldsymbol{p}_\perp$ is the hadron transverse momentum, with respect to the fragmenting quark direction $\boldsymbol{p}_q$. $\hat{\boldsymbol{p}}_q$ and $\hat{\boldsymbol{p}}_\perp$ denote unit vectors.

Eq. (7) shows that there might be a spin-$\boldsymbol{p}_\perp$ correlation in the fragmentation function, leading to an azimuthal asymmetry (for a unit transverse polarization vector, $S_q = 1$),

$$\begin{aligned} D_{h/q^\uparrow}(z, \boldsymbol{p}_\perp) - D_{h/q^\downarrow}(z, \boldsymbol{p}_\perp) &= \Delta^N\hat{D}_{h/q^\uparrow}(z, p_\perp)\,\boldsymbol{S}_q \cdot (\hat{\boldsymbol{p}}_q \times \hat{\boldsymbol{p}}_\perp) \\ &= \Delta^N\hat{D}_{h/q^\uparrow}(z, p_\perp)\,\sin(\phi_S - \phi_h)\,, \end{aligned} \tag{8}$$

where ϕ_S and ϕ_h are respectively the azimuthal angles, around the fragmenting quark direction, of the quark spin and the hadron transverse momentum. $\Delta^N\hat{D}_{h/q^\uparrow}$ is referred to as the Collins function; another common notation widely used in the literature, due to the Amsterdam group [10], is

$$\Delta^N\hat{D}_{h/q^\uparrow} = 2\,\frac{p_\perp}{zM_h}\,H_1^{\perp q}\,. \tag{9}$$



Figure 2: Three dimensional kinematics of the SIDIS process.

Such a basic quark asymmetry can originate an observable azimuthal asymmetry in polarized SIDIS processes, $\ell\, N^\uparrow \to \ell\, h\, X$, see Fig. 2 for kinematical configurations and notations. The virtual photon scatters off a transversely polarized quark inside the transversely polarized nucleon (in amounts described by h_1^q) and the quark polarization gets modified (it is reduced in size and tilted symmetrically with respect to the normal to the lepton plane, according to QED interactions). The final polarized quark then fragments according to Eq. (7). All this translates into a single spin asymmetry $(A_N^h)_{Coll}$:

$$(A_N^h)_{Coll} = \frac{\sum_q e_q^2 \int h_1^q(x)\,(1-y/(xy^2)\,\Delta^N\hat{D}_{h/q^\uparrow}(z,p_\perp)}{2\sum_q e_q^2 \int f_{q/p}(x)\,[1+(1-y)^2]/(xy^2)\,\hat{D}_{q/p}(z,p_\perp)}\,\sin(\phi_h+\phi_S) \tag{10}$$

which combines the transversity distribution and the Collins function. The integrals are usually performed in order to study the dependence of the asymmetry on one kinematical variable at a time, taking into account the experimental setups and the kinematical cuts. In the above expression we have neglected the intrinsic motion in the distribution functions; this also might originate azimuthal asymmmetries, for example with a $\sin(\phi_h - \phi_S)$ dependence, as we shall see in Subsection 2.1. Indeed, the numerator of the observed asymmetry,

$$A_N^h = \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow}\,, \tag{11}$$

contains several contributions. However, each of them has a different azimuthal dependence, so that one can extract the desired contribution by appropriately averaging A_N^h. The experimental data sensitive to the Collins mechanism, according to the so-called Trento conventions [11], are presented as:

$$A_{UT}^{\sin(\phi_h+\phi_S)} \equiv 2\,\frac{\int d\phi_h\, d\phi_S\, [d\sigma^\uparrow - d\sigma^\downarrow]\, \sin(\phi_h+\phi_S)}{\int d\phi_h\, d\phi_S\, [d\sigma^\uparrow + d\sigma^\downarrow]} \tag{12}$$

$$= \frac{\sum_q e_q^2 \int d\phi_h\, d\phi_S\, h_1^q(x)\, (1-y/(xy^2)\, \Delta^N\hat{D}_{h/q^\uparrow}(z,p_\perp)\, \sin^2(\phi_h+\phi_S)}{\sum_q e_q^2 \int d\phi_h\, d\phi_S\, f_{q/p}(x)\, [1+(1-y)^2]/(xy^2)\, \hat{D}_{q/p}(z,p_\perp)}\,.$$

Data on $A_{UT}^{\sin(\phi_h+\phi_S)}$ have been obtained by HERMES [12] and COMPASS [13] collaborations. The HERMES data and Eq. (12) have been used in Ref. [14], assuming a Soffer saturated value of $2|h_1^q(x)| = q(x) + \Delta q(x)$, to extract the Collins favoured, $\Delta^N\hat{D}_{\pi^+/u^\uparrow}$ and unfavoured, $\Delta^N\hat{D}_{\pi^+/d^\uparrow}$, fragmentation functions. It is the first attempted extraction of the Collins functions, which still have, given the quality and amount of the experimental information available, large uncertainties.

Furher information on the Collins function, which would greatly help in accessing the transversity distributions, is coming from BELLE collaboration. They can measure the combined effects of two Collins mechanisms by looking, event by event, at azimuthal correlations between hadrons in opposite jets in $e^+e^- \to h_1\, h_2\, X$ processes; first results [15] clearly indicate non zero Collins functions, in qualitative agreement with the Collins functions extracted from SIDIS data, as illustrated in Figs. 3 [16].

1.3 Alternative accesses to transversity

We have seen how the chiral-odd transversity distribution h_1^q must couple to another chiral-odd function in order to form a measurable quantity and to be experimentally accessible. The most promising approach seems to go via the double transverse spin asymmetry $A_{TT}^{p\bar{p}}$ in Drell-Yan processes with polarized protons *and antiprotons*; however, such an experiment is only at the proposal stage [6], with many difficulties yet to overcome [17]. At present, the most duable approach seems to be the azimuthal transverse SSA in SIDIS, in conjunction with data from BELLE. $A_{UT}^{\sin(\phi_h+\phi_S)}$ depends on the convolution of transversity and Collins functions; they are both unknown, but they depend on different variables; moreover, BELLE data should supply information on the product of two Collins functions alone.

Let us consider briefly other possible ways of coupling h_1^q to a chiral-odd function.

- *Inclusive Λ production and measurement of its polarization*



Figure 3: Simultaneous fit [16] of the HERMES data on $A_{UT}^{\sin(\phi_h+\phi_S)}$ [12], based on Eq. (12), (upper part) and of the BELLE data [15] on the convolution of two Collins functions (lower part). The two fits are equally good, showing consistency between the Collins functions extracted from SIDIS and e^+e^- processes. The resulting Collins functions are in qualitative agreement with those obtained in Ref. [14].

One should consider the electromagnetic SIDIS process $\ell\, p^\uparrow \to \gamma^* \to \ell\, \Lambda^\uparrow X$; the measurement of the Λ polarization $\boldsymbol{P}_\Lambda$, allows to study the spin transfer from the polarized quark to the polarized Λ. For transverse spins one has, schematically [18]

$$P_\Lambda \sim \sum_q h_1^q(x) \otimes \Delta_T D_{\Lambda/q}(z)\,, \tag{13}$$

which couples h_1^q to its (unknown) analogous fragmentation function, $\Delta_T D_{\Lambda/q} = D_{\Lambda^\uparrow/q^\uparrow} - D_{\Lambda^\downarrow/q^\uparrow}$. Notice that such an observable, a double spin asymmetry, does not vanish in the absence of intrinsic motions.

- *Two pion production and interference fragmentation functions*

It was suggested [19, 20], as an alternative to the Collins fragmentation mechanism, to consider the process $\ell\, p^\uparrow \to \ell\, \pi\, \pi\, X$, selecting two pions (or two hadrons) in the final state. In such a case, there might easily be a non zero chiral-odd, unknown, interference fragmentation function, δq_I. One should, like for the single pion Collins mechanism, look at a dependence on $\sin(\phi_R + \phi_S)$, where ϕ_R is the azimuthal angle of the relative momentum $\boldsymbol{R}$ of the two pions:

$$A_{UT} \sim \sum_q \sin(\phi_R + \phi_S)\, h_1^q \otimes \delta q_I. \tag{14}$$

This asymmetry has been observed by HERMES [21] and the interference fragmentation functions might be independently measured at BELLE.

- *Vector meson production in SIDIS*

  The off-diagonal element $\rho_{1,0}$ of the helicity density matrix of vector mesons produced in SIDIS processes is measurable and is related to h_1^q and some new (unknown) product of fragmentation amplitudes [18].

- *Inclusive hadronic production,* $p^\uparrow p \to \pi X$

  SSA in inclusive hadronic processes, $p^\uparrow p \to \pi X$, might depend on the convolution of transversity distributions and the Collins function. However, it has been shown recently that a careful treatment of the non collinear partonic interactions, strongly suppresses this contribution [22].

- *SSA in Drell-Yan processes*

  Data on unpolarized Drell-Yan processes show a strong $\cos(2\varphi)$ dependence, which might be explained [23] by the so-called Boer-Mulders function,

  $$\Delta^N f_{q^\uparrow/p} = -\frac{k_\perp}{M}\, f_1^{\perp q}\,, \tag{15}$$

  correlating the transverse spin of quarks inside an unpolarized proton to their transverse momentum $\boldsymbol{k}_\perp$ [10]. In such a case, even inside unpolarized protons one can find transversely polarized quarks and the elementary $q\,\bar{q} \to \ell^-\ell^+$ scattering of the Drell-Yan process shows the typical $\cos(2\varphi)$ dependence of double transverse spin asymmetries, Eq. (6). Then, one can learn about $\Delta^N f_{q^\uparrow/p}$.

  Drell-Yan processes may exhibit also single spin asymmetries, $d\sigma^\uparrow \neq d\sigma^\downarrow$, which originate from the convolution of transversity and the Boer-Mulders function:

  $$d\sigma^\uparrow - d\sigma^\downarrow \sim \sum_q h_1^q \otimes \Delta^N f_{q^\uparrow/p}. \tag{16}$$

  Thus, a combined study of the $\cos(2\varphi)$ dependence of unpolarized Drell-Yan processes (which yields information on $\Delta^N f_{q^\uparrow/p}$) and of transverse SSA, could allow to extract information on h_1^q [23, 24].

- *Transversity and GPD's*

  We briefly touch upon the role of transverse spin in the more general context of Generalized Parton Distributions (GPD), by mentioning that chiral-odd GPD's could be accessed in diffractive electroproduction of two vector mesons [25].

2. Transverse Single Spin Asymmetries (SSA)

Transverse Single Spin Asymmetries are intriguing obervables which test the underlying theory at a deep quantum mechanical level and, experimentally, very often result in unexpected and interesting data [26].

Let us consider, for example, the elastic scattering of two protons in their center of mass frame. If $\boldsymbol{p}$, $\boldsymbol{S}$ and $\boldsymbol{P}_T$ denote respectively the momentum of the incoming proton, its polarization vector and the transverse (with respect to the collision axis) momentum of the scattered proton, the only SSA allowed by parity invariance must be of the form:

$$A_S \equiv \frac{d\sigma^{\boldsymbol{S}} - d\sigma^{-\boldsymbol{S}}}{d\sigma^{\boldsymbol{S}} + d\sigma^{-\boldsymbol{S}}} \sim \boldsymbol{S} \cdot (\boldsymbol{p} \times \boldsymbol{P}_T) \sim P_T \sin(\phi_S - \phi) \,, \tag{17}$$

where ϕ_S and ϕ are the azimuthal angles, around the collision axis, of the spin vector and the scattered proton, respectively. Notice that A_S is zero if $\boldsymbol{S}$ lies in the scattering plane and if $P_T = 0$.

SSA are then possible in principle, with the azimuthal angular dependence of Eq. (17). Do they occurr at the partonic, pQCD, level? Let us consider the example of the elastic scattering of two identical quarks, in their center of mass frame, with the scattering taking place in the (xz) plane. All observables can be expressed in terms of 5 independent helicity amplitudes, $M_{\lambda'_1,\lambda'_2;\lambda_1,\lambda_2}$, which are usually denoted as:

$$\begin{array}{lll} M_{++;++} \equiv \Phi_1 & M_{--;++} \equiv \Phi_2 & M_{+-;+-} \equiv \Phi_3 \\ M_{-+;+-} \equiv \Phi_4 & M_{-+;++} \equiv \Phi_5 \,. & \end{array}$$

If only one of the two initial quarks is polarized along the y-direction with spin ↑ or ↓ and we do not look at the final quark polarizations, we can have $d\sigma^\uparrow \neq d\sigma^\downarrow$. In terms of the helicity amplitudes:

$$A_N \equiv \frac{d\sigma^\uparrow - d\sigma^\downarrow}{d\sigma^\uparrow + d\sigma^\downarrow} \sim \mathrm{Im}\,\left[\Phi_5\,\left(\Phi_1 + \Phi_3 + \Phi_2 - \Phi_4\right)^*\right] \,, \tag{18}$$

which can be non zero only if the single helicity flip amplitude Φ_5 is non zero and if different amplitudes have relative phases. The first condition implies a factor m_q/E_q (quark mass over its energy) and the second requires considering higher order contributions (at lowest order there are no relative phases). Altogether, one has

$$A_N^q \simeq \frac{m_q}{E_q}\,\alpha_s \,, \tag{19}$$

where α_s is the strong coupling constant. This makes single spin asymmetries in the partonic interactions entirely negligible [27]. However, sizeable SSA are observed at the hadronic level in elastic [28] and inclusive processes [29, 30, 31, 12].

2.1 SSA in SIDIS processes, data and extraction of the Sivers function

Transverse SSA can be understood, even within perturbative QCD or QED dynamics, by taking into account spin effects in the non perturbative components of the hard scattering factorized shemes. One such example has been discussed in Subsection 1.2, where the non perturbative spin-$\boldsymbol{p}_\perp$ dependent contribution is given by the Collins mechanism, Eq. (7), which leads to the SSA given in Eq. (10).

Another spin-intrinsic motion correlation can be introduced in the parton distribution functions, as originally suggested by Sivers [32]:

$$\hat{f}_{a/p^\uparrow}(x, \boldsymbol{k}_\perp) = \hat{f}_{a/p}(x, k_\perp) \ + \ \frac{1}{2}\,\Delta^N \hat{f}_{a/p^\uparrow}(x, k_\perp)\, \boldsymbol{S} \cdot (\hat{\boldsymbol{p}} \times \hat{\boldsymbol{k}}_\perp) \tag{20}$$

$$\hat{f}_{a/p^\uparrow}(x, \boldsymbol{k}_\perp) - \hat{f}_{a/p^\downarrow}(x, \boldsymbol{k}_\perp) \ = \ \Delta^N \hat{f}_{a/p^\uparrow}(x, k_\perp)\, \sin(\phi_S - \varphi)\,, \tag{21}$$

which shows how the number density of partons a, inside a transversely polarized proton with polarization $\boldsymbol{S}$, may depend on the relative azimuthal angle between $\boldsymbol{S}$ and the transverse momentum $\boldsymbol{k}_\perp$, $(\phi_S - \varphi)$.

In SIDIS processes, $\gamma^* p \to h X$ (Fig. 2.), the virtual photon interacts with a parton inside the transversely polarized proton, and the scattered parton then fragments into the observed hadron h, which has a transverse momentum $\boldsymbol{P}_T$, with respect to the γ^*-direction. This P_T can originate (at lowest order) from the initial intrinsic motion of the parton or from the fragmentation process. The latter case was considered in Subsection 1.2, and leads to a $\sin(\phi_h + \phi_S)$ dependence. The former case instead, explores the $\boldsymbol{k}_\perp$-distribution of partons inside a transversely polarized proton, and is then related to the Sivers function (21) $\Delta^N \hat{f}_{a/p^\uparrow} = -(2k_\perp/m_p) f_{1T}^{\perp q}$ [10]. A model for such a mechanism was introduced in Ref. [33].

The analogue of Eq. (10) now reads (see Ref. [34] for further details):

$$(A_N^h)_{Siv} \simeq \frac{\displaystyle\sum_q \int d^2\boldsymbol{k}_\perp\, \Delta^N f_{q/p^\uparrow}(x, k_\perp)\ \sin(\varphi - \phi_S)\, \frac{d\hat{\sigma}^{\ell q \to \ell q}}{dQ^2}\, D_{h/q}(z, p_\perp)}{\displaystyle 2\sum_q \int d^2\boldsymbol{k}_\perp\, f_{q/p}(x, k_\perp)\, \frac{d\hat{\sigma}^{\ell q \to \ell q}}{dQ^2}\, D_{h/q}(z, p_\perp)}\,, \tag{22}$$

where we have neglected some terms of order $(k_\perp^2/Q^2)$. Notice that, following Ref. [34], we have now taken into account all the intrinsic motion effects: in the distribution functions, in the elementary interactions

$$\frac{d\hat{\sigma}^{\ell q \to \ell q}}{dQ^2} = e_q^2\, \frac{2\pi\alpha^2}{\hat{s}^2}\, \frac{\hat{s}^2 + \hat{u}^2}{Q^4}\,, \tag{23}$$

and in the fragmentation functions.

This Sivers asymmetry is extracted from the data on A_N^h, Eq. (11), by selecting the $\sin(\phi_h - \phi_S)$ averaged asymmetry:

$$A_{UT}^{\sin(\phi_h-\phi_S)} \simeq \tag{24}$$

$$\frac{\displaystyle\sum_q \int d\phi_S\, d\phi_h\, d^2\boldsymbol{k}_\perp\; \Delta^N f_{q/p^\uparrow}(x,k_\perp)\sin(\varphi-\phi_S)\,\frac{d\hat\sigma^{\ell q\to \ell q}}{dQ^2}\; D_q^h(z,p_\perp)\sin(\phi_h-\phi_S)}{\displaystyle\sum_q \int d\phi_S\, d\phi_h\, d^2\boldsymbol{k}_\perp\; f_q(x,\boldsymbol{k}_\perp)\,\frac{d\hat\sigma^{\ell q\to \ell q}}{dQ^2}\; D_{h/q}(z,p_\perp)}\,.$$

Several groups [36] have exploited the above expression and the HERMES [12] and COMPASS [13] data to extract information, or check models, on the Sivers functions for u and d quarks. The fits to the $\pi^\pm$ HERMES data obtained in Ref. [35] are shown in Fig. 4, together with predictions for π^0 production.



Figure 4: HERMES data[12] on $A_{UT}^{\sin(\phi_\pi-\phi_S)}$ for scattering off a transversely polarized proton target and pion production; the curves are the results of the fit of Ref. [35]. The shaded area spans a region corresponding to one-sigma deviation at 90% CL. The curves for π^0 are predictions based on the extracted Sivers functions.

The resulting Sivers functions for u and d quarks turn out to be approximately opposite. This explains why the COMPASS data [13], taken on a deuterium target, show almost negligible values of $A_{UT}^{\sin(\phi_h-\phi_S)}$. Infact, for a deuterium target one has:

$$A_{UT}^{\sin(\phi_h-\phi_S)} \sim \left(\Delta^N f_{u/p^\uparrow} + \Delta^N f_{d/p^\uparrow}\right)\left(4\, D_{h/u} + D_{h/d}\right)\;. \tag{25}$$

2.2 SSA in $p^\uparrow p \to \pi\, X$ processes

Let us conclude by mentioning SSA in $p^\uparrow p \to \pi\, X$ processes. Some recent papers have discussed the problem, in the context of QCD with a possible factorization scheme [37] and/or with higher-twist partonic correlations [38].

We only mention here that both Sivers and Collins mechanism, assuming a QCD factorization scheme with parton intrinsic motions, might contribute to a non vanishing A_N; however, it was recently shown that the correct treatment of the elementary dynamics, with non collinear partonic processes and the proper spinor phases taken into account, strongly suppresses the contribution of the Collins mechanism [22]. The Sivers mechanism, instead, is not suppressed [39] and can well explain the observed SSA [30, 31]; the Sivers functions active in $\ell\, p$ and $p\, p$ inclusive processes might be the same.

Acknowledgements

I would like to acknowledge invaluable help from my collaborators M. Boglione, U. D'Alesio, A. Kotzinian, E. Leader, S. Melis, F. Murgia and A. Prokudin. This work is partially supported by the European Community – Research Infrastructure Activity under the FP6 "Structuring the European Research Area" programme (HadronPhysics, contract number RII3-CT-2004-506078).

References

[1] See, *e.g.*: V. Barone, A. Drago and P. Ratcliffe, *Phys. Rep.* **359** (2002) 1

[2] J. Ralston and D.E. Soper, *Nucl. Phys.* **B152** (1979) 109; J.L. Cortes, B. Pire and J.P. Ralston, *Zeit. Phys.* **C55** (1992) 409; R.L. Jaffe and X. Ji, *Nucl. Phys.* **B375** (1992) 527

[3] J.C. Collins, *Nucl. Phys.* **B396** (1993) 161

[4] V. Barone, T. Calarco and A. Drago, *Phys. Rev.* **D56** (1997) 527

[5] O. Martin, A. Schäfer, M. Stratmann and W. Vogelsang, *Phys. Rev.* **D57** (1998) 3084; *Phys. Rev.* **D60**, (1999) 117502

[6] PAX Collaboration, e-Print Archive: hep-ex/0505054

[7] M. Anselmino, V. Barone, A. Drago and N. N. Nikolaev, *Phys. Lett.* **B594** (2004) 97

[8] A.V. Efremov, K. Goeke and P. Schweitzer, *Eur. Phys. J.* **C35**, (2004) 207

[9] H. Shimizu, G. Sterman, W. Vogelsang and H. Yokoya, *Phys. Rev.* **D71** (2005) 114007

[10] P.J. Mulders and R.D. Tangerman, *Nucl. Phys.* **B461** (1996) 197; *Erratum-ibid.* **B484** (1997) 538; D. Boer and P.J. Mulders, *Phys. Rev.* **D57** (1998) 5780; D. Boer, P.J. Mulders and F. Pijlman, *Nucl. Phys.* **B667** (2003) 201

[11] A. Bacchetta, U. D'Alesio, M. Diehl and C.A. Miller, *Phys. Rev.* **D70** (2004) 117504

[12] HERMES Collaboration, A. Airapetian *et al.*, *Phys. Rev. Lett.* **94** (2005) 012002; M. Diefenthaler, e-Print Archive: hep-ex/0507013

[13] COMPASS Collaboration, V.Yu. Alexakhin *et al.*, *Phys. Rev. Lett.* **94** (2005) 202002

[14] W. Vogelsang and F. Yuan, *Phys.Rev.* **D72** (2005) 054028

[15] Belle Collaboration (K. Abe *et al.*), e-Print Archive: hep-ex/0507063

[16] A. Prokudin, private communication

[17] N.N. Nikolaev and F.F. Pavlov, e-Print Archive: hep-ph/0512051

[18] M. Anselmino, M. Boglione, J. Hansson and F. Murgia, *Phys. Rev.* **D54** (1996) 828

[19] R.L. Jaffe, X. Jin and J. Tang, *Phys. Rev. Lett.* **80**, (1998) 1166

[20] M. Radici, R. Jakob and A. Bianconi, *Phys. Rev.* **D65**, (2002) 074031

[21] P. van der Nat, talk delivered on behalf of the HERMES collaboration at Transversity 2005, Villa Olmo (Como), 7–10th. September 2005

[22] M. Anselmino, M. Boglione, U. D'Alesio E. Leader and F. Murgia, *Phys. Rev.* **D71** (2005) 014002

[23] D. Boer, *Phys. Rev.* **D60** (1999) 014012

[24] A. Bianconi and M. Radici, *J. Phys.* **G31** (2005) 645

[25] D.Yu. Ivanov, B. Pire, L. Szymanowski, O.V. Teryaev, *Phys. Lett.* **B550** (2002) 65

[26] For a recent review, see A.D. Krisch, e-Print Archive: hep-ex/0511040

[27] G.L. Kane, J. Pumplin and W. Repko, *Phys. Rev. Lett.* **41**, (1978) 1689

[28] D.G. Crabb *et al.*, *Phys. Rev. Lett.* **65** (1990) 3241

[29] K. Heller *et al.*, *Phys. Rev. Lett.* **51** (1983) 2025

[30] D.L. Adams *et al.* (E704 Collaboration), *Z. Phys.* **C56**, (1992) 181; *Phys. Lett.* **B345** (1995) 569

[31] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92** (2004) 171801

[32] D. Sivers, *Phys. Rev.* **D41** (1990) 83; **D43** (1991) 261

[33] S.J. Brodsky, D.S. Hwang and I. Schmidt, *PHys. Lett.* **B530**, (2002) 99

[34] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, *Phys. Rev.* **D71** (2005) 074006

[35] M. Anselmino, M. Boglione, U. D'Alesio, A. Kotzinian, F. Murgia and A. Prokudin, *Phys. Rev.* **D72** (2005) 094007

[36] M. Anselmino *et al.*, e-Print Archive: hep-ph/0511017

[37] M. Anselmino, M. Boglione, U. D'Alesio, E. Leader, S. Melis and F. Murgia, e-Print Archive: hep-ph/0509035

[38] A. Bacchetta, C.J. Bomhof, P.J. Mulders and F. Pijlman, *Phys. Rev.* **D72** (2005) 034030; A. Bacchetta, e-Print Archive: hep-ph/0511085

[39] U. D'Alesio and F. Murgia, *Phys. Rev.* **D70** (2004) 074009